

# AQUINAS COLLEGE HEALTH AND IMMUNIZATION RECORD

**PART I**

must be turned in prior to beginning of semester

**PAGE 1 OF 6**

Name \_\_\_\_\_  
First Name Middle Name

---

Last Name

---

Address \_\_\_\_\_  
Street City State Zip

---

Date of Entry    /   /         Date of Birth    /   /         School ID# \_\_\_\_\_  
M Y      M D Y

---

Status:    Part-time \_\_\_\_\_    Full-time \_\_\_\_\_    Graduate \_\_\_\_\_    Undergraduate \_\_\_\_\_    Professional \_\_\_\_\_

**PART II: TO BE COMPLETED AND SIGNED BY YOUR HEALTH CARE PROVIDER.**

*All information must be in English.*

**A. MMR (MEASLES, MUMPS, RUBELLA)**

(Two doses required at least 28 days apart for students born after 1956 and all health care professional students.)

1. Dose 1 given at age 12 months or later ..... #1    /   /     
M D Y
2. Dose 2 given at least 28 days after first dose ..... #2    /   /     
M D Y

**B. MENINGOCOCCAL QUADRIVALENT**

(A, C, Y, W-135) One or 2 doses for all college students; revaccinate every 5 years if increased risk continues.

1. Quadrivalent conjugate (preferred; administer simultaneously with Tdap if possible).

a. Dose #1    /   /         b. Dose #2    /   /     
M D Y      M D Y

2. Quadrivalent polysaccharide (acceptable alternative if conjugate not available).

Date    /   /     
M D Y

**C. TETANUS, DIPHTHERIA, PERTUSSIS**

1. Primary series completed?    Yes     No       Date of last dose in series:    /   /     
M D Y

2. Date of most recent booster dose:    /   /         Type of booster:    Td     Tdap   
M D Y      *Tdap booster recommended for ages 11-64 unless contraindicated*

**D. HEPATITIS B**

(All college and health care professional students. Three doses of vaccine or two doses of adult vaccine in adolescents 11–15 years of age, or a positive hepatitis B surface antibody meets the requirement.)

1. Immunization (hepatitis B)

a. Dose #1    /   /         b. Dose #2    /   /         c. Dose #3    /   /     
M D Y      M D Y      M D Y

Adult formulation     Child formulation       Adult formulation     Child formulation       Adult formulation     Child formulation

2. Immunization (Combined hepatitis A and B vaccine)

a. Dose #1    /   /         b. Dose #2    /   /         c. Dose #3    /   /     
M D Y      M D Y      M D Y

3. Hepatitis B surface antibody    Date    /   /         Result:    Reactive     Non-reactive   
M D Y

# AQUINAS HEALTH AND IMMUNIZATION RECORD (CONTD.)

## E. INFLUENZA

Trivalent (IIV3) \_\_\_\_\_ Quadrivalent (IIV4) \_\_\_\_\_ Recombinant (RIV3) \_\_\_\_\_ Live attenuated influenza vaccine (LAIV) \_\_\_\_\_

Date of last dose: \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y

## F. VARICELLA

(Birth in the U.S. before 1980, a history of chicken pox, a positive varicella antibody, or two doses of vaccine meets the requirement.)

1. History of Disease Yes \_\_\_ No \_\_\_ or Birth in U.S. before 1980 Yes \_\_\_ No \_\_\_

2. Varicella antibody \_\_\_\_/\_\_\_\_/\_\_\_\_ Result: Reactive \_\_\_\_\_ Non-reactive \_\_\_\_\_  
M D Y

3. Immunization

a. Dose #1 ..... #1 \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y

b. Dose #2 given at least 12 weeks after first dose ages 1–12 years..... #2 \_\_\_\_/\_\_\_\_/\_\_\_\_  
and at least 4 weeks after first dose if age 13 years or older. M D Y

## G. HUMAN PAPILLOMAVIRUS VACCINE (HPV2/HPV4/HPV9)

(Three doses of vaccine for females and males 11–26 years of age at 0, 1–2, and 6 month intervals.)

Immunization (indicate which preparation, if known) Quadrivalent (HPV4) \_\_\_\_\_ or Bivalent (HPV2) \_\_\_\_\_ or 9-valent (HPV9) \_\_\_\_\_

a. Dose #1 \_\_\_\_/\_\_\_\_/\_\_\_\_ b. Dose #2 \_\_\_\_/\_\_\_\_/\_\_\_\_ c. Dose #3 \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y M D Y

## H. HEPATITIS A

1. Immunization (hepatitis A)

a. Dose #1 \_\_\_\_/\_\_\_\_/\_\_\_\_ b. Dose #2 \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y

2. Immunization (Combined hepatitis A and B vaccine)

a. Dose #1 \_\_\_\_/\_\_\_\_/\_\_\_\_ b. Dose #2 \_\_\_\_/\_\_\_\_/\_\_\_\_ c. Dose #3 \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y M D Y

## I. PNEUMOCOCCAL POLYSACCHARIDE VACCINE

PCV 13 \_\_\_\_\_ Date \_\_\_\_/\_\_\_\_/\_\_\_\_ PPSV 23 \_\_\_\_\_ Date \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y

## J. MENINGOCOCCAL SEROUGROUP B

(Two or three dose series; may be given to any college student or for outbreak control; may be given with quadrivalent meningococcal vaccine at different anatomic site. Must complete series with the same vaccine.)

1. MenB-RC (Bexsero) \_\_\_ routine \_\_\_ outbreak –related

a. Dose #1 \_\_\_\_/\_\_\_\_/\_\_\_\_ b. Dose #2. \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y

OR

1. MenB-FHbp (Trumenba) \_\_\_ routine \_\_\_ outbreak-related

a. Dose #1 \_\_\_\_/\_\_\_\_/\_\_\_\_ b. Dose #2 \_\_\_\_/\_\_\_\_/\_\_\_\_ c. Dose #3 \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y M D Y

## I. POLIO

(Primary series, doses at least 28 days apart. Three primary series are acceptable. See ACIP website for details.)

1. OPV alone (oral Sabin three doses): #1 \_\_\_\_/\_\_\_\_/\_\_\_\_ #2 \_\_\_\_/\_\_\_\_/\_\_\_\_ #3 \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y M D Y

2. IPV/OPV sequential: IPV #1 \_\_\_\_/\_\_\_\_/\_\_\_\_ IPV #2 \_\_\_\_/\_\_\_\_/\_\_\_\_ OPV #3 \_\_\_\_/\_\_\_\_/\_\_\_\_ OPV #4 \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y M D Y M D Y

3. IPV alone (injected Salk four doses): #1 \_\_\_\_/\_\_\_\_/\_\_\_\_ #2 \_\_\_\_/\_\_\_\_/\_\_\_\_ #3 \_\_\_\_/\_\_\_\_/\_\_\_\_ #4 \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y M D Y M D Y

# AQUINAS HEALTH AND IMMUNIZATION RECORD (CONTD.)

## M. TUBERCULOSIS (TB) SCREENING/TESTING<sup>1</sup>

Please answer the following questions:

Have you ever had close contact with persons known or suspected to have active TB disease?  Yes  No

Were you born in one of the countries or territories listed below that have a high incidence of active TB disease?  Yes  No  
(If yes, please CIRCLE the country, below)

Afghanistan	Congo	Iran (Islamic Republic of)	Namibia	Solomon Islands
Algeria	Côte d'Ivoire	Iraq	Nauru	Somalia South Africa
Angola	Democratic People's Republic of Korea	Kazakhstan	Nepal	South Sudan
Anguilla	Democratic Republic of the Congo	Kenya	Nicaragua	Sri Lanka
Argentina	Djibouti	Kiribati	Niger	Sudan
Armenia	Dominican Republic	Kuwait	Nigeria	Suriname
Azerbaijan	Ecuador	Kyrgyzstan	Northern Mariana Islands	Swaziland
Bangladesh	El Salvador	Lao People's Democratic Republic	Pakistan	Tajikistan
Belarus	Equatorial Guinea	Latvia	Palau	Thailand
Belize	Eritrea	Lesotho	Panama	Timor-Leste
Benin	Estonia	Liberia	Papua New Guinea	Togo
Bhutan	Ethiopia	Libya	Paraguay	Trinidad and Tobago
Bolivia (Plurinational State of)	Fiji	Lithuania	Peru	Tunisia
Bosnia and Herzegovina	French Polynesia	Madagascar	Philippines	Turkmenistan
Botswana	Gabon	Malawi	Poland	Tuvalu
Brazil	Gambia	Malaysia	Portugal	Uganda
Brunei Darussalam	Georgia	Maldives	Qatar	Ukraine
Bulgaria	Ghana	Mali	Republic of Korea	United Republic of Tanzania
Burkina Faso	Greenland	Marshall Islands	Republic of Moldova	Uruguay
Burundi	Guam	Mauritania	Romania	Uzbekistan
Cabo Verde	Guatemala	Mauritius	Russian Federation	Vanuatu
Cambodia	Guinea	Mexico	Rwanda	Venezuela (Bolivarian Republic of)
Cameroon	Guinea-Bissau	Micronesia (Federated States of)	Saint Vincent and the Grenadines	Viet Nam
Central African Republic	Guyana	Mongolia	Sao Tome and Principe	Yemen
Chad	Haiti	Montenegro	Senegal	Zambia
China	Honduras	Morocco	Serbia	Zimbabwe
China, Hong Kong SAR	India	Mozambique	Seychelles	
China, Macao SAR	Indonesia	Myanmar	Sierra Leone	
Colombia			Singapore	
Comoros				

Source: World Health Organization Global Health Observatory, Tuberculosis Incidence 2014. Countries and territories with incidence rates of  $\geq 20$  cases per 100,000 population. For future updates, refer to <http://www.who.int/tb/country/en/>.

Have you had frequent or prolonged visits\* to one or more of the countries or territories listed above with a high prevalence of TB disease? (If yes, CHECK the countries or territories, above)  Yes  No

Have you been a resident and/or employee of high-risk congregate settings (e.g., correctional facilities, long-term care facilities, and homeless shelters)?  Yes  No

Have you been a volunteer or health care worker who served clients who are at increased risk for active TB disease?  Yes  No

Have you ever been a member of any of the following groups that may have an increased incidence of latent *M. tuberculosis* infection or active TB disease: medically underserved, low-income, or abusing drugs or alcohol?  Yes  No

**If the answer is YES to any of the above questions**, Aquinas College requires that you receive TB testing as soon as possible but at least prior to the start of the subsequent semester).

**If the answer to all of the above questions is NO**, no further testing or further action is required.

\* The significance of the travel exposure should be discussed with a health care provider and evaluated.

# AQUINAS HEALTH AND IMMUNIZATION RECORD (CONTD.)

### TUBERCULOSIS (TB) RISK ASSESSMENT (to be completed by **health care provider**)

Clinicians should review and verify the information above. Persons answering YES to any of the questions in Part M are candidates for either Mantoux tuberculin skin test (TST) or Interferon Gamma Release Assay (IGRA), unless a previous positive test has been documented.

History of a positive TB skin test or IGRA blood test? (If yes, document below) Yes \_\_\_\_\_ No \_\_\_\_\_

History of BCG vaccination? (If yes, consider IGRA if possible.) Yes \_\_\_\_\_ No \_\_\_\_\_

#### 1. TB Symptom Check

Does the student have signs or symptoms of active pulmonary tuberculosis disease? Yes \_\_\_\_\_ No \_\_\_\_\_ *If No, proceed to 2 or 3*

If yes, check below:

- Cough (especially if lasting for 3 weeks or longer) with or without sputum production
- Coughing up blood (hemoptysis)
- Chest pain
- Loss of appetite
- Unexplained weight loss
- Night sweats
- Fever

Proceed with additional evaluation to exclude active tuberculosis disease including tuberculin skin testing, chest x-ray, and sputum evaluation as indicated.

#### 2. Tuberculin Skin Test (TST)

(TST result should be recorded as actual millimeters (mm) of induration, transverse diameter; if no induration, write “0”. The TST interpretation should be based on mm of induration as well as risk factors.)\*\*

Date Given: \_\_\_\_/\_\_\_\_/\_\_\_\_ Date Read: \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y

Result: \_\_\_\_\_ mm of induration \*\*Interpretation: positive\_\_\_\_ negative\_\_\_\_

Date Given: \_\_\_\_/\_\_\_\_/\_\_\_\_ Date Read: \_\_\_\_/\_\_\_\_/\_\_\_\_  
M D Y M D Y

Result: \_\_\_\_\_ mm of induration \*\*Interpretation: positive\_\_\_\_ negative\_\_\_\_

#### \*\*Interpretation guidelines

>5 mm is positive:

- Recent close contacts of an individual with infectious TB
- persons with fibrotic changes on a prior chest x-ray, consistent with past TB disease
- organ transplant recipients and other immunosuppressed persons (including receiving equivalent of >15 mg/d of prednisone for >1 month.)
- HIV-infected persons

>10 mm is positive:

- recent arrivals to the U.S. (<5 years) from high prevalence areas or who resided in one for a significant\* amount of time
- injection drug users
- mycobacteriology laboratory personnel
- residents, employees, or volunteers in high-risk congregate settings
- persons with medical conditions that increase the risk of progression to TB disease including silicosis, diabetes mellitus, chronic renal failure, certain types of cancer (leukemias and lymphomas, cancers of the head, neck, or lung), gastrectomy or jejunoileal bypass and weight loss of at least 10% below ideal body weight.

>15 mm is positive:

- persons with no known risk factors for TB who, except for certain testing programs required by law or regulation, would otherwise not be tested.

\* The significance of the travel exposure should be discussed with a health care provider and evaluated.

#### 3. Interferon Gamma Release Assay (IGRA)

Date Obtained: \_\_\_\_/\_\_\_\_/\_\_\_\_ (specify method) QFT-GIT T-Spot other\_\_\_\_  
M D Y

Result: negative\_\_\_\_ positive\_\_\_\_ indeterminate\_\_\_\_ borderline\_\_\_\_ (T-Spot only)

Date Obtained: \_\_\_\_/\_\_\_\_/\_\_\_\_ (specify method) QFT-GIT T-Spot other\_\_\_\_  
M D Y

Result: negative\_\_\_\_ positive\_\_\_\_ indeterminate\_\_\_\_ borderline\_\_\_\_ (T-Spot only)

# AQUINAS HEALTH AND IMMUNIZATION RECORD (CONTD.)

## 4. Chest x-ray: (Required if TST or IGRA is positive)

Date of chest x-ray: \_\_\_/\_\_\_/\_\_\_ Result: normal \_\_\_ abnormal \_\_\_  
M D Y

### Management of Positive TST or IGRA

All students with a positive TST or IGRA with no signs of active disease on chest x-ray should receive a recommendation to be treated for latent TB with appropriate medication. However, students in the following groups are at increased risk of progression from LTBI to TB disease and should be prioritized to begin treatment as soon as possible.

- Infected with HIV
- Recently infected with *M. tuberculosis* (within the past 2 years)
- History of untreated or inadequately treated TB disease, including persons with fibrotic changes on chest radiograph consistent with prior TB disease
- Receiving immunosuppressive therapy such as tumor necrosis factor-alpha (TNF) antagonists, systemic corticosteroids equivalent to/greater than 15 mg of prednisone per day, or immunosuppressive drug therapy following organ transplantation
- Diagnosed with silicosis, diabetes mellitus, chronic renal failure, leukemia, or cancer of the head, neck, or lung
- Have had a gastrectomy or jejunioileal bypass
- Weigh less than 90% of their ideal body weight
- Cigarette smokers and persons who abuse drugs and/or alcohol

••Populations defined locally as having an increased incidence of disease due to *M. tuberculosis*, including medically underserved, low-income populations

\_\_\_\_\_ Student agrees to receive treatment

\_\_\_\_\_ Student declines treatment at this time

---

### PART III. HEALTH FORM

Parent's or guardian's name \_\_\_\_\_ Phone (\_\_\_\_) \_\_\_\_\_

Father's place of employment \_\_\_\_\_ Phone (\_\_\_\_) \_\_\_\_\_

Mother's place of employment \_\_\_\_\_ Phone (\_\_\_\_) \_\_\_\_\_

Family physician \_\_\_\_\_ Phone (\_\_\_\_) \_\_\_\_\_

Health insurance policy information. Insurance carrier \_\_\_\_\_

Contract No. \_\_\_\_\_

Group No. \_\_\_\_\_

Subscriber's name \_\_\_\_\_ Street Address \_\_\_\_\_

City, State, Zip \_\_\_\_\_ Telephone (\_\_\_\_) \_\_\_\_\_

Subscriber's DOB \_\_\_\_\_

---

---

# AQUINAS HEALTH AND IMMUNIZATION RECORD (CONTD.)

## PART IV. MEDICAL HISTORY

Current medical/psychological concerns \_\_\_\_\_  
\_\_\_\_\_

Past medical/psychological concerns \_\_\_\_\_  
\_\_\_\_\_

Surgeries/Hospitalizations \_\_\_\_\_  
\_\_\_\_\_

Drug allergies: \_\_\_\_\_

Other Allergies: \_\_\_\_\_

Current medications and dosage \_\_\_\_\_  
\_\_\_\_\_

Recommendations and/or restrictions \_\_\_\_\_  
\_\_\_\_\_

\*\*\*\*\*If on medication for ADHD or ADD, please provide documentation from prescribing physician.\*\*\*\*\*

## HEALTH CARE PROVIDER

Name \_\_\_\_\_ Signature \_\_\_\_\_

Address \_\_\_\_\_ Phone (\_\_\_\_\_) \_\_\_\_\_

The above is for informational use only. I understand that my personal information may be shared with the Aquinas College Student Affairs Department with the purpose of assisting me in achieving academic success and continuity of care.

Student signature \_\_\_\_\_ Date \_\_\_\_\_

Parent/Guardian signature \_\_\_\_\_ Date \_\_\_\_\_

Please send completed form to:

**Aquinas College**  
**Counseling, Health & Wellness Services, Lower Donnelly Center**  
**1700 Fulton Street E, Grand Rapids, MI 49506**  
**or CHWS@aquinas.edu**  
**Questions: Counseling, Health & Wellness Services, 616-632-2905.**

**AQUINAS COLLEGE**